



AGENCE DES NORMES ET DE LA QUALITÉ®
STANDARDS AND QUALITY AGENCY®

PROJET DE NORME CAMEROUNAISE PNC 3155 : 2026 - CT 48

2026

www.anor.cm

Cassava and cassava products — Determination of total cyanogens — Enzymatic assay method

Toute reproduction ou représentation intégrale ou partielle, par quelque procédé que ce soit, des pages publiées dans le présent document, faite sans l'autorisation de l'éditeur est illicite et constitue une contrefaçon. Seules sont autorisées d'une part, les reproductions strictement réservées à l'usage privé du copiste et non destinées à une utilisation collective et, d'autre part, les analyses et courtes citations justifiées par le caractère scientifique ou d'information de l'œuvre dans laquelle elles sont incorporées.

PROJET DE NORME CAMEROUNAISE

ENQUETE PUBLIQUE N° : 19

Durée de l'enquête Du 21/09/2026 Au 19/10/2026

Edition et diffusion par l'Agence des Normes et de la Qualité

B.P.: 14966 Yaoundé – CAMEROUN – Tél: 6992 257 777 /Fax.: (237) 222 22 64 96

E-mail : enquetepublique@anor.cm – www.anor.cm/enquetes-publiques

ANOR®

DN-R2/PR-01/ENR-02
16/12/2015

**Cassava and cassava products — Determination of total
cyanogens — Enzymatic assay method**



Table of contents

1	Scope	1
2	Normative references	1
3	Principle.....	1
4	Reagents	1
5	Apparatus	2
6	Procedure	2
6.1	Sample preparation	2
6.2	Analytical assay	2
6.3	Colorimetric procedure	2
6.4	Calculation of cyanogen content	2
7	Test report	3
	Annex A (normative) Determination of moisture content	4
	Bibliography	5
	Acknowledgment and appreciation	5

Foreword

The African Organization for Standardization (ARSO) is an African intergovernmental organization established by the United Nations Economic Commission for Africa (UNECA) and the Organization of African Unity (AU) in 1977. One of the fundamental mandates of ARSO is to develop and harmonize African Standards (ARS) for the purpose of enhancing Africa's internal trading capacity, increase Africa's product and service competitiveness globally and uplift the welfare of African communities. The work of preparing African Standards is normally carried out through ARSO technical committees. Each Member State interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, Regional Economic Communities (RECs), governmental and non-governmental organizations, in liaison with ARSO, also take part in the work.

ARSO Standards are drafted in accordance with the rules given in the ISO/IEC Directives, Part 2.

The main task of technical committees is to prepare ARSO Standards. Draft ARSO Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an ARSO Standard requires approval by at least 75 % of the member bodies casting a vote.

Attention is drawn to the possibility that some of the elements of this document may be the subject of patent rights. ARSO shall not be held responsible for identifying any or all such patent rights.

This African Standard was prepared by the ARSO Technical Harmonization Committee on *Agriculture and Food Products* (ARSO/THC 02).

© African Organisation for Standardisation 2016 — All rights reserved*

ARSO Central Secretariat
International House 3rd Floor
P. O. Box 57363 — 00200 City Square
NAIROBI, KENYA

Tel. +254-20-2224561, +254-20-3311641, +254-20-3311608

E-mail: arso@arso-oran.org

Web: www.arso-oran.org

* © 2016 ARSO — All rights of exploitation reserved worldwide for African Member States' NSBs.

Copyright notice

This ARSO document is copyright-protected by ARSO. While the reproduction of this document by participants in the ARSO standards development process is permitted without prior permission from ARSO, neither this document nor any extract from it may be reproduced, stored or transmitted in any form for any other purpose without prior written permission from ARSO.

Requests for permission to reproduce this document for the purpose of selling it should be addressed as shown below or to ARSO's member body in the country of the requester:

© African Organisation for Standardisation 2016 — All rights reserved

ARSO Central Secretariat
International House 3rd Floor
P.O. Box 57363 — 00200 City Square
NAIROBI, KENYA

Tel: +254-20-2224561, +254-20-3311641, +254-20-3311608

E-mail: arso@arso-oran.org
Web: www.arso-oran.org

Reproduction for sales purposes may be subject to royalty payments or a licensing agreement. Violators may be prosecuted.

Introduction

Cassava (*Manihot esculenta* Crantz) is one of the staple food crops. A major drawback of cassava utilization is its potential toxicity due to the presence of endogenous cyanogenic glucosides.

Processing of cassava should reduce the cyanogenic glucosides to an acceptable level. This test method therefore, is intended to be used for determination of the cyanogen content in cassava and its different products as a measure to safeguard the health of the consumers.

ENQUÊTE PUBLIQUE N° 19

Cassava and cassava products — Determination of total cyanogens — Enzymatic assay method

1 Scope

This African Standard specifies a method for the determination of total cyanogens in cassava and cassava products.

2 Normative references

The following referenced documents are indispensable for the application of this document. For dated references, only the edition cited applies. For undated references, the latest edition of the referenced document (including any amendments) applies.

No normative references have been cited in this standard.

3 Principle

Linamarase hydrolyses cyanogenic glucoside compounds of cassava extract in orthophosphoric acid medium to produce cyanohydrins which rapidly decompose to cyanide ion in alkali (NaOH), followed by addition of excess pH 6 buffer and chloramine-T to produce a purple solution which is measured spectrophotometrically

4 Reagents

4.1 Linamarin, stock solution (5 mmol) made by dissolving 31.0 mg of Linamarin in 25 ml buffer, pH 6.

NOTE Weighing should be done quickly as the compound is hygroscopic.

4.2 Linamarase, stock solution made by dissolving linamarase enzyme in a 0.1 M sodium phosphate buffer to obtain a 2.5 µg linamarin in 0.1 M aliquot

4.3 Isonicotinic acid

4.4 1,3-dimethyl barbituric acid

4.5 Acetone cyanohydrin (99 %), stock (5 mmol) made by first dissolving 570 µl (0.532 g) in 25ml of 0.1 M orthophosphoric acid, and from this solution, making 1.00 ml up to 50 ml with 0.1 M orthophosphoric acid.

Because of the poor performance of the pure compound in a pipette, the added quantity should also be weighed for accurate calculation of concentration. The refrigerated stocks can be stored for several weeks to months.

4.6 Potassium cyanide (KCN), stock (5 mmol) made by dissolving 163 mg dry and pure KCN in 500 ml 0.2 M NaOH. KCN should be dried for at least 12 h over concentrated H₂SO₄.

Standard solutions of 80 µmol and 320 µmol are made from the stock solutions by making 1.60 ml of them up to, respectively 100 ml and 25 ml with 0.1 M orthophosphoric acid. This results in 8 nmol/tube and 32 nmol/tube and absorbance of about 0.220 absorbance unit (AU) and 0.880 AU depending on the spectrophotometer. KCN standards are made just prior to analysis and applied as 0.1 ml to 0.5 ml orthophosphoric acid and 3.4 ml buffer, pH 6 or as 0.1 ml to 3.9 ml buffer, pH 4.

4.7 Chloramine T solution.

4.8 Colour reagent, pure NaOH (3.7 g) is dissolved in 200 ml distilled water. 7 g of 1,3-dimethyl barbituric acid and 5.7 g isonicotinic acid are dissolved in this alkaline solution by extensive stirring. The pH is adjusted between 7 and 8 with 1M HCl or NaOH. This reagent can be kept for at least 12 days at room temperature.

NOTE Stock solutions and reagents should be of analytical grade.

5 Apparatus

Ordinary laboratory apparatus, and a spectrophotometer

6 Procedure

6.1 Sample preparation

Cut fresh cassava root or moist products into 1 cm cubes and randomly take 50 g cassava cubes and homogenize in 250 ml refrigerated 0.1M orthophosphoric acid in a blender for 15 s at low speed, followed by 60 s at high speed, 60 s of rest and 60 s at full speed again.

For flour 4 g is swirled gently in 25 ml of refrigerated extraction medium in a 50 ml closed container for 5 min.

Extracts are best analyzed immediately, but if not, they should be stored refrigerated or frozen for up to two months.

6.2 Analytical assay

6.2.1 Cyanogenic potential.

0.1 ml extract is added to 0.4 ml buffer, pH 7, in a test tube, followed by addition of 0.1 ml linamarase solution. After 15 min incubation at 30 °C, 0.6 ml NaOH (0.2 M) is added, followed after 5 min, by 2.8 ml buffer, pH 6. Coloration is as described in 6.3.

6.2.2 Non glucosidic cyanogens

0.1 ml extract is added to 0.6 ml of NaOH (0.2 M). After 2 min, 3.3 ml buffer, pH 6 is added, followed by coloration (6.3).

6.2.3 Free cyanide (HCN)

0.6 ml extract is diluted with 3.4 ml buffer, pH 6 and assayed by colorimetry (6.3).

NOTE Cyanogens are assayed in duplicate and all assays are carried out in glass-stoppered test tubes. The contents of the tubes are mixed after each addition.

6.3 Colorimetric procedure

0.1 ml Chloramine-T reagent (2 % w/v) is added to the 4 ml buffered extract in the test tubes and mixed. After 5 min, 0.6 ml of color reagent is added and mixed. The absorbance at 605 nm is measured after 10 min. Reagent blanks are run for each analysis. In a digital single beam spectrophotometer, the absorbance can be measured between 0.050 AU and 2.000 AU (Absorbance Unit). In less sophisticated analog equipment, absorbance can be measured between 0.100 AU and 1.200 AU with acceptable accuracy.

6.4 Calculation of cyanogen content

Cyanogen levels are calculated in mg HCN equivalent per kilogram sample on dry weight basis (mg HCN equivalent kg⁻¹, DWB) as follows:

$$\text{HCN equivalent} = \frac{0.027 \chi \left[v + \frac{sm}{100} \right]}{ds \left[1 - \frac{m}{100} \right]}$$

where,

s is the sample weight in grams (g);

v is the volume, in millilitres, of the extraction medium;

d is the volume, in millilitres, of extract assayed; and

m is the moisture content, %.

Dry 5 g to 10 g sample (to constant weight), at 105 °C ± 2 °C (oven) or 70 °C in a vacuum oven and calculate the moisture content (%).

χ is the quantity of cyanogens (nmol) in the tube, calculated from calibration curve as follows:

$$\chi = \frac{(A_{605} - w)}{\text{Slope}}$$

where,

A_{605} is the absorbance measured at 605 nm. Both slope and intercept "w" are derived by linear regression of the calibration points by means of a calculator. For very low values of "w" (<0.005) and higher values of A_{605} (>0.200) the intercept can be neglected.

7 Test report

The test report shall show the method used and the result obtained.

It shall also mention any operating conditions not specified in this standard, or regarded as optional, as well as any circumstances that may have influenced the result.

The report shall include all details required for the complete identification of the sample analyzed.

Annex A
(normative)

Determination of moisture content

A.1 Procedure

Weigh accurately 10 g of the material in a suitable moisture dish previously dried in an electric oven and weighed. Place the dish in an electric oven maintained at $105\text{ }^{\circ}\text{C} \pm 1\text{ }^{\circ}\text{C}$ for 5 h. Cool the dish in a desiccator and weigh with the lid on. Repeat the process of heating, cooling and weighing at half-hour intervals until the loss in weight between two successive weighings is less than 1 mg. Record the lowest weight obtained.

A.2 Calculation and expression of results

$$\text{Moisture, percent by mass} = \frac{(M_1 - M_2) \times 100}{M_1 - M_3}$$

where,

M_1 is the mass, in grams, of the dish and sample before drying;

M_2 is the mass, in grams, of the dish and sample after drying;

M_3 is the mass, in grams, of the dish only.

Bibliography

EAS 744:2010, *Cassava and cassava products — Determination of total cyanogens — Enzymatic assay method*

Acknowledgment and appreciation

This African Standard was harmonized under the project titled:

Harmonization of African Standards for Agriculture and Food Products
June 2015 — August 2018

financed by the **African Development Bank (AfDB)** through the **African Trade Fund (AfTra)** Programme. ARSO acknowledges and appreciates this support on behalf of its Member States.

ENQUÊTE PUBLIQUE N° 19

ENQUETE PUBLIQUE N° 19